



|            |   |
|------------|---|
| Title      | 銀河系内に於ける星の運動について  |
| Author(s)  | 福田, 守   |
| Citation   | 北海道學藝大學紀要. 第二部, 8(2): 11-13   |
| Issue Date | 1957-12   |
| URL        | <a href="http://s-ir.sap.hokkyodai.ac.jp/dspace/handle/123456789/5544">http://s-ir.sap.hokkyodai.ac.jp/dspace/handle/123456789/5544</a> |
| Rights     |   |

## On the Motions of Stars in the Galaxy

Mamoru FUKUDA

Department of Astronomy, Hokkaido Gakugei  
University, Asahikawa

福田 守： 銀河系内に於ける星の運動について

### Abstract

As a model of the galaxy a non-homogeneous spheroid is considered whose equidensity surfaces are similar and similarly situated spheroids. For density distribution we use the sum of two normal distribution functions whose parameters were determined so as to fit a recent radio observation. The coefficients in the expressions for stellar motions in the galaxy are calculated.

### § 1 Introduction

Many observations<sup>1)</sup> indicate that the isophotal curves of an elliptical nebula are ellipses having a common centre and almost the same axis ratio. We therefore assume that the equidensity surfaces of the galaxy are similar ellipsoids.

In 1956, Contopoulos<sup>2)</sup> solved the equations of motion of a star in the non-homogeneous spheroid with an arbitrary mass distribution.

On the other hand, several models<sup>3)</sup> have been considered to represent the mass distribution of the galaxy. In 1955, Takase<sup>4)</sup> assumed a model of the normal density distribution, and determined the parameters so as this model to represent a recent radio observation.

### § 2 The potential function

Let the equidensity ellipses be similar to a basic ellipsoid with axes  $a$ ,  $b$ , and  $c$ . If  $a_1$ ,  $b_1$ , and  $c_1$  are the axes of any equidensity ellipsoid, then

$$\frac{a_1}{a} = \frac{b_1}{b} = \frac{c_1}{c} = \sqrt{u}.$$

A potential per unit mass at an interior point  $P(x, y, z)$  of a non-homogeneous ellipsoid with density function  $\sigma(u)$  is

$$U = -\pi Gabc \int_0^\infty \frac{d\lambda}{D} \int_\infty^u \sigma(u) du$$

where

$$D^2 = (a^2 + \lambda)(b^2 + \lambda)(c^2 + \lambda)$$

and

$$u = \frac{x^2}{a^2 + \lambda} + \frac{y^2}{b^2 + \lambda} + \frac{z^2}{c^2 + \lambda}.$$

If  $\sigma(u)$  can be expanded in a Taylor series we have

$$\sigma(u) = \sigma_0 + \sigma_1 u + \cdots + \frac{\sigma_n}{n!} u^n + \cdots \quad (1)$$

Now we consider the spheroid ( $a=b>c$ ), and then

$$u = \frac{r^2}{a^2 + \lambda} + \frac{z^2}{c^2 + \lambda}$$

and

$$U - U_0 = -\pi G \left[ \sigma_0 \left\{ A_{10}(r^2 - r_0^2) + A_{01}(z^2 - z_0^2) \right\} + \cdots \right. \\ \left. + \frac{\sigma_{n-1}}{n!} \sum \frac{n!}{\mu! \nu!} A_{\mu\nu} (r^{2\mu} z^{2\nu} - r_0^{2\mu} z_0^{2\nu}) + \cdots \right]$$

where  $\mu + \nu = n$  and

$$A_{\mu\nu} = a^2 c \int_0^\infty \frac{d\lambda}{(a^2 + \lambda)^{\mu+1} (c^2 + \lambda)^{\nu+1/2}}.$$

The values of  $A_{\mu\nu}$  for various values of  $c/a$  can be calculated.<sup>2), 5)</sup>

### § 3 Equations of motion

We consider the spheroidal system, and the projection of the attracting force on the plane of symmetry is radial. Therefore, using cylindrical co-ordinates  $r$ ,  $\theta$  and  $z$ , we have

$$r^2 \frac{d\theta}{dt} = C \quad (2)$$

$$\frac{C^2}{r^2} + \left( \frac{dr}{dt} \right)^2 + \left( \frac{dz}{dt} \right)^2 = 2U(r, z) + H \quad (3)$$

$$\frac{d^2 z}{dt^2} = \frac{\partial U}{\partial z}. \quad (4)$$

The solution of system (3) and (4) can be obtained by the method of undetermined coefficients.

Let  $r$  and  $z$  be given by the series

$$\left. \begin{aligned} r &= r_0 + r_1 t + r_2 t^2 + \cdots \\ z &= z_0 + z_1 t + z_2 t^2 + \cdots \end{aligned} \right\} \quad (5)$$

Whenever  $z_0 = 0$ , we also have  $z_2 = 0$ , and from (3), (4), and (5) we find

$$r = r_0 + r_1 t + \left( \frac{C^2}{2r_0^3} - \pi G \sigma_0 A_{10} r_0 + \cdots \right) t^2 + \cdots \quad (6)$$

and

$$z = z_1 t - \frac{\pi G}{3} z_1 (\sigma_0 A_{01} + \sigma_1 A_{11} r_0^2 + \cdots) t^3 + \cdots \quad (7)$$

### § 4 Density distribution

TAKASE<sup>4)</sup> assumed following density distribution function

$$\rho(r) = \rho_1(r) + \rho_2(r) = \rho_1(0) e^{-h_1^2 r^2} + \rho_2(0) e^{-h_2^2 r^2} \quad (8)$$

and determined the parameters so as the circular rotational velocity derived from this

model to represent the observational result by Kwee, Muller and Westerhout,<sup>6)</sup> who gave the rotational velocity curve over the wide range extending to the inner part of the galaxy from their 21cm observation.

Assuming  $c/a=1/12$  he obtained following values :

$$\begin{aligned} h_1 &= 0.19 kpc^{-1} & M_1 &= 0.591 \times 10^{11} M_o \\ h_2 &= 0.60 kpc^{-1} & M_2 &= 0.089 \times 10^{11} M_o. \end{aligned}$$

According to the relation<sup>4)</sup>

$$M_i = \frac{\sqrt{\pi^3}}{h_i^3} \frac{\sqrt{1-\varepsilon^2}}{\rho_i(O)}$$

we find

$$\begin{aligned} \rho_1(O) &= 1.88 \times 10^8 M_o kpc^{-3} \\ \rho_2(O) &= 8.92 \times 10^8 M_o kpc^{-3}. \end{aligned}$$

### § 5 Calculations of coefficients

Inserting  $r^3 = a^2 u$  into the right hand side of (8) and developpe those exponential functions, we have

$$\rho(u) = \rho_1(O) + \rho_2(O) - a^2 \{ h_1^2 \rho_1(O) + h_2^2 \rho_2(O) \} u + \dots \quad (9)$$

From (1) and (9) we find

$$\begin{aligned} \sigma_0 &= \rho_1(O) + \rho_2(O) = 1.08 \times 10^9 M_o kpc^{-3} \\ \sigma_1 &= -a^2 \{ h_1^2 \rho_1(O) + h_2^2 \rho_2(O) \} = -7.38 \times 10^{10} M_o kpc^{-3}. \end{aligned}$$

Furthermore we have  $A_{10}=0.13$ ,  $A_{01}=2.00^5)$ ,  $a^2 A_{11}=1.67^2)$ ,  $a=15 kpc$  and  $C=2.83 \times 10^6$ . Inserting these values into (6) and (7) we obtain

$$\begin{aligned} r &= r_0 + r_1 t + (4.00 r_0^{-3} - 2.95 \times 10^{-14} r_0) t^2 + \dots \\ z &= z_1 t - 7.00 \times 10^{-23} z_1 (2.16 \times 10^9 - 5.84 \times 10^8 r_0^2) t^3 + \dots \end{aligned}$$

Giving  $r_0$ ,  $r_1$ , and  $z_1$  we can calculate the motion of the star. If we find the solution  $r=r(t)$  and  $z=z(t)$ ,  $\theta$  is given by the following formula

$$\theta - \theta_0 = \int_{r_0}^r \frac{C dt(r)}{r^2}.$$

### References

- 1) For example, J. H. Oort, Ap. J. **91**, 273, 1940.
- 2) G. Contopoulos, Ap. J. **124**, 643, 1956.
- 3) For example, J. H. Oort, B. A. N. **9**, 185, 1941.
- 4) B. Takase, Publ. A. S. Japan, **7**, 201, 1955.
- 5) H. Mineur, Ann. d'Ap., **2**, 1, 1939.
- 6) K. K. Kwee, C. A. Muller and G. Westerhout, B. A. N., **12**, 211, 1954.